

Synthesis of Some Mannich Base Derivatives and their Antifungal, Antibacterial Activities Study

Dr.K. S. Manjunatha

Department of Pharmaceutical Chemistry, Kuvempu University, Post Graduate Centre, Kadur-577548, Karnataka, India.

Abstract

Some new Mannich bases (**1a-5a**) were synthesized by reacting morpholine with various aldehydes and the compounds containing active hydrogen atoms (acetophenone, thiourea and phenylurea). Structures of the compounds were confirmed by IR, ¹H-NMR, ¹³C-NMR and mass spectra. Antimicrobial studies were carried out for the synthesized compounds against various bacterial and fungal strains employing Ofloxacin as standard drug for antibacterial and amphotericin B as standard drug for antifungal screening. The compounds **1a-3a** and **5a** have been possessed high antibacterial activity and the compounds **3a** and **5a** showed higher zone of inhibition against fungal strains. Particularly the compound **3a** exhibited higher zone of inhibition against *A. niger* and the compound **5a** showed higher activity against *A. niger* and *Trichophyton*.


Keywords: Mannich condensation; Antibacterial activity; Antifungal activity

1. Introduction

In recent years serious attention has been directed towards the discovery and development of new antimicrobial drugs. The main problem in the treatment of fungal infection is the increasing drug resistance subjected to antimycotic therapy such as persons infected with HIV [1]. Nosocomial infections during the past decade have invaded hospitals worldwide by multidrug resistant Gram-positive and Gram-negative pathogens. Searching for novel antimicrobial agents and new microbial targets is in demand to intervene to avert the danger caused by these life-threatening infections. Since the resistance towards the available antibiotics among pathogenic bacteria has grown rapidly, there is a clear need for the development of new and effective antimicrobial agents. Therefore, the success in designing antimicrobial agents which are distinct from those of the classical antibiotics is the key for treating such infectious diseases known for their chronicity and failure to treat with conventional antibiotics which will eventually lead to death [2,3].

Mannich bases have gained importance due to their applications in pharmaceutical industry and other applications including agrochemicals and plant growth regulators. The presence of heterocyclic moieties in organic compounds is of interest in biology, pharmacology, optics, materials science and so on [4-7]. The chemistry of the aminoalkylation of aromatic substrates by the Mannich reaction is of great interest for the synthesis and modification of biologically active compounds having physical and chemical importance as well as physiological properties because the amino group can be easily converted into a variety of other functionalities [8]. Mannich reaction offers a judicious method for introduction of basic aminoalkyl chain in various drugs/compounds. Further a considerable amount of work has been reported on synthesis and pharmacological activities as well as intermediates in drug synthesis [9]. Morpholine derivatives were reported to possess antimicrobial, anti-inflammatory and central nervous system activities [10-12].

Some of the Mannich bases containing heterocyclic ring and their names are given below. Therefore, bearing in mind the above observation, we were led to synthesize new series of Mannich base derivatives.



2. Materials and methods

All the reagents and solvents for synthesis were purchased from Alfa Aesar and Sigma Aldrich Chemicals. Silicagel-GF coated glass plates were used for monitoring the reactions by iodine vapors as visualizing agents and methanol and acetone solvent system were used as TLC mobile phase. Melting points of compounds were determined using digital melting point apparatus and were uncorrected. The IR spectra were obtained on SHIMADZU FTIR spectrophotometer (KBr pellets, λ -max in cm^{-1}) and $^1\text{H-NMR}$ spectra of compounds were recorded on Bruker Model-300 using DMSO-*d*₆ as solvent. Mass spectra of the synthesized compounds were recorded on Jeol SX-102/DA-6000 spectrometer.

2.1. General procedure for the synthesis of the compounds (1a-5a)

To an ethanolic solution of thiourea (0.76g, 0.01M), 3 pellets of sodium hydroxide and morpholine (0.8mL, 0.01M) were added followed by anisaldehyde (1.2mL, 0.01M). The reaction mixture was taken in a RB flask and kept over a magnetic stirrer. The mixture was stirred under ice cold condition for 8h. The precipitate formed was filtered, washed with water, dried and recrystallized from methanol. The reaction scheme is shown in figure 1.

2.2. Scheme





Figure 1-Scheme of the reaction



3. Results and Discussion

1-((4-methoxyphenyl)(morpholino)methyl)thiourea (1a).

IR (cm^{-1}) KBr disc: 3343.24-NH str; 2960-CH str; 1605-NH bend $^1\text{H-NMR}$ (ppm): 9.81 Hof NH; 7.82 Hof-NH₂; 7.3-6.94 Hof Aromatic-H; 5.61 Hof-CH; 3.8-3.48 Hof morpholine $^{13}\text{C-NMR}$ (ppm): 191 C of C=S; 164-1136 C of aromatic-C; 661 C of-CH; 61-554 C of morpholine m/e: 281.

1-(morpholino(thiophen-2-yl)methyl)-3-phenylurea (2a).

IR (cm^{-1}) KBr disc: 3295-NH str; 2960-CH str; 1637.90 C=O str; 1560-NH bend $^1\text{H-NMR}$ (ppm): 8.51 Hof NH; 7.5-7.08 Hof of Aromatic-H; 6.91 Hof NH; 5.81 Hof CH; 3.3-3.28 Hof Morpholine $^{13}\text{C-NMR}$ (ppm): 154 C=O, 144, 139, 128, 126, 124, 121, 117 aromatic-C, 68, 66 (Morpholine), 47 (CH) m/e: 317.

3-morpholino-1-phenyl-3-(thiophen-2-yl)propan-1-one(3a).

IR(cm⁻¹)**KBr disc**: 3459–NHstr; 3064–CHstr; 1653C=Ostr; 1591–NHbend¹**H-NMR**(ppm): 8.0-6.98Hof AromaticH; 4.81HofCH; 2.92HofCH₂; 3.5-2.58HofMorpholine¹³**C-NMR**(ppm): 205C=O, 145, 143, 139, 131, 128, 127, 126, 125, 124, 123aromatic-C, 74(CH₂), 51(CH), 38, 36(Morpholine)**m/e**: 301.

1-(morpholino(pyridin-2-yl)methyl)-3-phenylurea(4a).

IR(cm⁻¹)**KBr disc**: 3425–NHstr; 3035–CHstr; 1655CHstr; 1594–NHbend¹**H-NMR**(ppm): 9.21HofNH; 8.51HofAromaticH; 6.41HofNH; 5.81HofCH; 3.3-2.58Hofmorpholine¹³**C-NMR**(ppm): 1571CofCO; 154-11711CofaromaticC; 801CofCH; 62-594Cofmorpholinem/e: 312.

3-morpholino-1-phenyl-3-(pyridin-2-yl)propan-1-one(5a).

IR(cm⁻¹)**KBr disc**: 3412–NHstr; 2938–CHstr; 1678C=Ostr; 1587–NHbend¹**H-NMR**(ppm): 8.41Hof AromaticH; 7.9-7.48HofaromaticH; 7.21HofCH; 4.4-3.912HofCH₂; 3.5-2.38Hofmorpholinem/e: 297.

The compound **1a** was synthesized using anisaldehyde, Morpholine and acetophenone. Compounds **2a** and **3a** were prepared by reacting thiophene-2-carboldehyde, morpholine and active hydrogen compound such as phenylurea/acetophenone. Compound **4a** and **5a** were synthesized by adding Morpholine, pyridine-2-carboldehyde and active hydrogen compound such as phenylurea/acetophenone.

The molecular formula, colour, yield and melting point of the compounds are given in the Table 1. Melting points of the synthesized compounds were sharp indicating that the compounds are pure. The yield of the compounds is also suggested that the chemical methods were reliable for the synthesis of the compound. All spectral data were found to be match with the proposed structures.

Table.1: Analytical data of the synthesized compounds

S.No.	Molecular Formula	Molecular Weight	Melting Point (°C)	Colour	Yield %
1a	C ₁₃ H ₁₉ N ₃ O ₂ S	281.135	Yellow solid	90	
2a	C ₁₆ H ₁₉ NO ₂ S	317.144	White solid	95	
3a	C ₁₇ H ₁₉ NO ₂ S	301.175	White Solid	90	
4a	C ₁₇ H ₂₀ N ₄ O ₂	312.134	Brown Solid	92	
5a	C ₁₈ H ₂₀ N ₂ O ₂	296.90	Brown Solid	98	

3.2 Antimicrobial activity

The synthesized compounds were subjected for antimicrobial activity and the results were shown in the table 2. Three bacterial strains, two gram positive and one gram negative namely *B. subtilis*, *Enterobacter* and *Moraxella* were taken for antibacterial screening and two fungal pathogens, *A. niger* and *Trichophyton* were tested for antifungal studies against the synthesized compounds using Ofloxacin and amphotericin B as standard drug. All the compounds, except **4a** are highly active against *B. subtilis*. The compound **5a** exhibits good activity against *Enterobacter* whereas the rest of the compounds show only lesser activity. The increased bacterial activity against *Moraxella* was exhibited by the compounds **5a** and **1a**. The activity of other compounds against *Moraxella* is found to be very less. All the compounds are active against the fungi, *A. niger*. The compounds **3a** and **5a** possessed higher activity and others showed lesser activity. The compound **5a** exhibits higher zone of inhibition against *Trichophyton* whereas the other compounds showed lower activity and the compound **4a** showed no activity. The structural features of the compounds like presence of phenyl ring, the heterocyclic compounds containing nitrogen and electron donating group like -OCH₃ group were the cause for increased biological activity. While other compounds which possess electron withdrawing substituent responsible for lesser activity. The enhanced antibacterial activity of the compound **1a** is due to the presence of thiocarbonyl sulfur (C=S) [13]. In case of antifungal activity, there are several reports available that the derivatives with a N-containing heterocycles [14] and the compounds which contain acetophenone show significant activity [15-17].




Figure 2: Antimicrobial screening of the synthesized compounds

4. Conclusion

Five new Mannich bases were synthesized incorporating different aldehydes with various active hydrogen compounds and morpholine. Synthesized compounds were characterized by FTIR, ^1H & ^{13}C -NMR and mass spectral studies. These derivatives were evaluated for their antimicrobial activity against three bacterial strains including two gram positive and one gram negative bacteria and two fungal strains. Almost all the compounds possessed good activity against bacteria and fungal strains. Compound **1a** showed higher zone of inhibition against *B. subtilis* due to the presence of C=S group. The compound **5a** showed higher antifungal activity compared to the standard drug. Pyridine ring is responsible for the stronger antifungal activity.

References

- [1] Idhayadhulla, R. Surendra Kumar, A. Jamal Abdul Nasser, J. Selvin, Aseer Manilal, Synthesis of some Mannich base derivatives and their antimicrobial activity study, *Arabian Journal of Chemistry* 2014; 7(6).
- [2] Robert C. Spencer, Bacteremia caused by multi-resistant Gram-positive microorganisms, *Clinical Microbiology and Infection* 1999; 5(Supplement 2).
- [3] Suzan A. Matar, Wamidh H. Talib, Mohammad S. Mustafa, Mohammad S. Mubarak, Murad A. Al-Damen, Synthesis, characterization, and antimicrobial activity of Schiff bases derived from benzaldehydes and 3,3'-diaminodipropylamine, *Arabian Journal of Chemistry* 2013; 17.
- [4] Marcus Baumann, Ian R. Baxendale, Steven V. Ley and Nikzad Nikbin, An overview of the key routes to the best-selling 5-membered ring heterocyclic pharmaceuticals *Beilstein J. Org. Chem.* 2011.
- [5] Amir M, Kumar Hand Javed SA, Condensed bridgehead nitrogen heterocyclic system: synthesis and pharmacological activities of 1,2,4-triazolo-[3,4-b]-1,3,4-thiadiazole derivatives of ibuprofen and biphenyl-4-yloxyacetic acid, *Eur J Med Chem.* 2008.
- [6] Fatma A. Bassyouni, Tamer S. Saleh, Mahmoud M. El Hefnawi, Sherein I. Abd El-Moez, Waled M. El-Senousy, Mohamed E. Abdel-Rehim, Synthesis, pharmacological activity evaluation and molecular modeling of new polynuclear heterocyclic compounds containing benzimidazole derivatives, *Archives of Pharmacol Research* 2012; 35(12).

- [7] Suha K Al-Mosawi, Synthesis and Characterization of Heterocyclic Schiff Base, Thiazolidinone and Chalcone as Antibacterial Agents, *Research Journal of Pharmaceutical, Biological and Chemical Sciences* ISSN:0975-8585.
- [8] Idhayadhulla A., Surendra Kumar R., Jamal Abdul Nasser A., Selvin, Aseer Manilal J., Synthesis of some Mannich base derivatives and their antimicrobial activity study, *Arabian Journal of Chemistry* 2014; 7(6).
- [9] Sheela Joshi, Purti Bilgaiyan, Anju Pathak, Synthesis, spectroscopic characterization and antibacterial screening of novel Mannich bases of Ganciclovir *Arabian Journal of Chemistry* 5 March 2013.
- [10] P. Panneerselvam, M. Gnanarupa Priya, N. Ramesh Kumar, and G. Saravanan Synthesis and Pharmacological Evaluation of Schiff Bases of 4-(2-Aminophenyl)-Morpholines *Indian J Pharm Sci.* 2009 Jul-Aug.
- [11] Hakan Bektaş, Şule Ceylan, Neslihan Demirbaş, Şengül Alpay - Karaoğlu, Bahar Bilgin Sökmen, Antimicrobial and antiurease activities of newly synthesized morpholine derivatives containing an azole nucleus *Medicinal Chemistry Research*, Nov 2012.
- [12] J. Thanusu, V. Kanagarajan, M. Gopalakrishnan, Synthesis and Spectral Analysis of An Array of Novel 4-(4-morpholinophenyl)-6-aryl-pyrimidin-2-amines *J. Chil. Chem. Soc.* 2010; 55.
- [13] Ryukou Tokuyama, Yoshiei Takahashi, Yayoi Tomita, Tomio Suzuki, Toshihiko Yoshida, Nobuhiko Iwasaki, Noriyuki Kado, Eiichi Okezaki, and Osamu Nagata, Structure-Activity Relationship (SAR) Studies on Oxazolidinone Antibacterial Agents *Chem. Pharm. Bull.* 2001; 49(4) 347-352.
- [14] Hao Wang, Shuang-Xi Ren, Ze-Yu He, De-Long Wang, Xiao-Nan Yan, Jun-Tao Feng and Xing Zhang, Synthesis, Antifungal Activities and Qualitative Structure Activity Relationship of Carabrone Hydrazone Derivatives as Potential Antifungal Agents *Int. J. Mol. Sci.* 2014, 15(3), 4257-4272; doi: 10.3390
- [15] Ertan R, Ayhan G, Yulug N, Rolland Y, Synthesis and antifungal activities of some new triazolyl acetophenone derivatives *Arzneimittelforschung*. 1995 May.
- [16] Ma YT, Fan HF, Gao YQ, Li H, Zhang AL, Gao JM, Natural products as sources of new fungicides (I): synthesis and antifungal activity of acetophenone derivatives against phytopathogenic fungi. *Chem Biol Drug Des.* 2013 Apr.
- [17] Amar Patil, Rahul Jadhav, Hemant Raundal, Lokesh Sharma, Rupali Badgujar and Vivek Bobade, Synthesis and antifungal activities of diaryl pyrazole carboxamide derivatives. *Journal of Chemical and Pharmaceutical Research*, 2014, 6(8): 218-223.